

# PRELIMINARY CHECKLIST OF ENSIFERA (ORTHOPTERA) SPECIES FROM MT. MALINDANG RANGE NATURAL PARK, MINDANAO, PHILIPPINES

Rodrin R. Rivera<sup>1,2</sup>, Olga M. Nuñez<sup>1</sup>, Johniel Banquil<sup>1</sup>, Mark G. Robson<sup>3</sup>, Ming Kai Tan<sup>4</sup>

<sup>1</sup>Mindanao State University – Iligan Institute of Technology, Iligan City

<sup>2</sup>Northwestern Mindanao State College of Science and Technology

<sup>3</sup>Rutgers University – The State University of New Jersey, USA

<sup>4</sup>Lee Kong Chian Natural History Museum, National University of Singapore, Singapore

Volume 10, Issue No.1

## Abstract

Mt. Malindang Range Natural Park (MMRNP) is a key biodiversity area that serves as a crucial habitat for Ensifera species. This study provides the first preliminary checklist of Ensifera species in Mt. Malindang. The study was conducted through a purposive sampling via hand catching across 14 selected sites in MMRNP from January to November 2024 and from March to April 2025 respectively. Results showed a total of 36 species from six families, 11 subfamilies, and 28 genera of Ensifera were recorded with 25% endemic to Mindanao Island. *Paranisitra flavofacia* Baroga-Barbecho, Yap, Tan & Robillard, 2019 was the most dominant species recorded in the area. Interestingly, *Anthracites zebra* Hebard, 1922, was noted to have color variations, from which each species was collected at different sites. Additionally, two new species were identified in this study such *Anthracites seneaureus* sp. nov. Tan, Rivera & Daguplo, 2025 and *Scytocera musa* sp. nov. Tan & Rivera, 2025. The study highlights the underappreciated status of Ensiferans in Mt. Malindang, evidenced by the numerous new locality records and the identification of two previously unknown species. These results underscore the importance of ongoing and thorough exploration of Ensifera species in the area, building on sampling efforts that have been ongoing since 1959. Nonetheless, the lack of molecular data limits a complete understanding of their genetic composition and evolutionary connections, hindering efforts to accurately resolve phylogenetic relationships, assess cryptic diversity, and implement informed conservation strategies.

**Keywords:** Ensiferan checklist, Mt. Malindang, Species, Taxonomy

## INTRODUCTION

The Philippines is one of the megadiverse countries

worldwide, characterized by complex and rich biodiversity hotspots (Koh et al., 2019). With more than 7,000 islands, many

of which are considered species-rich and are left unexplored (Heads, 2013), the country harbors a remarkable diversity of insects, including orthopterans (crickets, grasshoppers, and katydids), which play a vital role in ecological processes. Orthopterans, with more than 2,000 known species across Southeast Asia (Tan et al., 2017a), contribute to nutrient cycling, soil aeration, and trophic interactions as both herbivores and prey for higher taxa (Tan et al., 2013). However, due to the archipelagic nature and vast ecological gradients of the Philippine islands, many habitats remain insufficiently surveyed, leaving numerous orthopteran species undiscovered (Conservation International Philippines, 2025; Robillard & Yap, 2015).

The order Orthoptera is classified into two major suborders: Caelifera, which includes short-horned grasshoppers, and Ensifera, which comprises long-horned grasshoppers, crickets, and katydids (Shin et al., 2024). The last comprehensive survey of the orthopteran checklist in the Philippines was compiled by Bruner over a century ago (Bruner, 1915). The compilation of a catalogue from this work highlighted the documentation

of various species of grasshoppers, crickets, and katydids, which served as a foundational reference for subsequent studies on Philippine Orthoptera. Moreover, since its publication, the archipelago's orthopteran diversity has been further explored, leading to the discovery and description of new species (Tan et al., 2017). For instance, a 2019 survey on Siargao Island (Tan et al., 2019) discovered 37 species and 34 genera emphasizes the rich orthopteran biodiversity in understudied areas. Tan et al. (2024) discovered a new genus and species *Fartmanntettix undulatus*, Tan & Tummbrinck, 2024 under subfamily Cladonotinae (Orthoptera: Tetrigidae) from Mindanao Island, tentatively classified under the tribe Trusmaditetrigini Storozhenko, 2023. This newly identified genus is distinguished from other members of the tribe by its undulating and elevated curinae on the pronotum and femora. Patano et al. (2021) described *Tegotettix derijeii* (Orthoptera: Tetrigidae), a newly identified pygmy grasshopper species from Mindanao. Previously known only from historical descriptions and illustrations, this species had not been documented for over a century. Meanwhile, Ensifera – a notable group of orthoptera are

recognized for their unique and diverse communication strategies such as acoustic signaling that is crucial for their mating and territorial behaviors (Walker & Rentz, 2020). This group of insects occupy a variety of microhabitats from forest canopies, leaf litters, and beneath the ground and serve as essential components of food webs both as herbivores and prey for larger animals (Guido & Gianelle, 2001). A recent expedition on Ensifera species in the Philippines, particularly in Mindanao, resulted in a taxonomic review of the genus *Paranisitra* and *Lebinthus*. This study led to the description of new species and new locality records of Ensifera from the different regions of Mindanao, highlighting the ongoing need for taxonomic revisions and biodiversity assessments in the region (Baroga-Barbecho et al., 2019; 2020). Additionally, Tan et al. (2023) conducted a bioacoustics study on the little-known Grylloidea crickets from various areas in Mindanao which led to the discovery of four previously undocumented species such as *Mistshenkoana lunotan*, *Mistshenkoana higaonon*, and *Ectatoderus dubius* from Misamis Oriental; and *Phaloria* (*Phaloria*) *rotundata* from Agusan del Sur, Philippines re-

spectively. Further, Bahoy et al. (2024) discovered new species, *Falcerminthus hispidus* of subtribe *Lebinthina* (Grylloidea, Eneopterinae, *Lebebinthini*) from Zamboanga del Norte and a new locality record of *Lebinthus luae* in Lake Sebu, Cotabato City in Mindanao, Philippines.

The Mt. Malindang Range Natural Park (MMRNP) is an ASEAN heritage park situated in the province of Misamis Occidental, Philippines. The park extends a total of 53,262 hectares of forest lands with covering grasslands to mossy-montane forest measuring at about 34,000 hectares that straddle the provinces of Misamis Occidental and Zamboanga Peninsula (DENR-BMB, 2016). The unique structure of Mt. Malindang is crucial as it is designated as one of the major terrestrial key biodiversity areas (KBA) in the country which provides environmental habitats for orthoptera specifically, Ensifera – a key indicator for maintaining ecological balance (Calago et al., 2019). The first attempt to sample the Ensifera species in the area was carried out by Tan et al., (2024) and discovered new species (*Itara mindanao*) from the subgenus *Itara* (Orthoptera: Grylloidea: Itarinae). The recent discovery was a first record of

the subgenus in the Philippines and in Mindanao Island, offering more insights on ensiferan biodiversity in the mountain range (Tan et al., 2024). This suggests that despite their presence, Ensifera species are often overlooked and largely unexplored. Given the ecological and economic significance of MMRNP as a major hub for economic activities and a designated key biodiversity area (Conservation International Philippines et al., 2006), documenting the Ensifera species and their distribution is essential for understanding and conserving the park's biodiversity.

This study presents the first preliminary checklist of Ensifera species recorded in Mt. Malindang Range Natural Park (MMRNP), accompanied by taxonomic notes. Additionally, it provides distribution data and potential biogeographic patterns for each species, along with detailed illustrations to aid in accurate species identification.

## **MATERIALS AND METHOD**

The study was conducted across 14 selected sites, encompassing 14 barangays within two cities and six municipalities in Misamis Occidental, all within the protected landscapes of

MMRNP which exhibit a distinctive geographical configuration characterized by bushy to mossy forest, dipterocarp forest, grasslands, and agricultural areas. The study took place from January to November 2024 and from March to April 2025 respectively. The northern portion of MMRNP comprises higher elevation (colder region) extending from 1600 to 2,400 m.a.s.l. The area is also characterized by a crater lake, waterfalls (Municipality of Don Victoriano) and hot springs (Oroquiety City) that paved a way for microhabitat composition for various species of Ensifera. The southern portion of MMRNP comprises areas with lower elevations and are characterized by riparian systems, vegetation covers, and crops.

Prior to sampling, the researcher secured a Wildlife Gratuitous Permit (GWP No. R10-2024-17) from the Department of Environment and Natural Resources – Region 10 (DENR-10). In addition, local permits were obtained from the relevant local government units (LGUs), barangays, and Indigenous Peoples through the Free, Prior, and Informed Consent (FPIC) process, ensuring the legal and ethical compliance of data collection.

Ensiferan specimens were collected using opportunistic sampling methods, including hand-catching and sweep-netting, conducted during both daytime (0600–1000 hours) and nighttime (1730–2230 hours) across multiple visits (at least three per site) from January to November 2024. Manual collection of adult Ensifera was carried out using wide-mouthed plastic vials, with most specimens located and captured during nighttime calling activity, although early morning collections were also made.


Specimens were preserved in 70% ethanol for later analysis. Acoustic recordings of calling songs were obtained by placing individuals in mesh cages and recording with a Zoom H1n recorder set at a 120 Hz sampling frequency. Recordings were saved in 24-bit, BWF-compliant WAV format. In situ photographs of the Ensiferan specimens were taken using iPhone XR and iPhone 15 smartphones.

The terminology and analytical procedures for acoustic data followed Tan et al. (2023). Temporal parameters, such as call durations and intervals, were measured manually using Raven Lite 2.0.0. Peak fre-

quency was analyzed using the *'spectro\_analysis'* function from the R package warbleR (Araya-Salas & Smith-Vidaurre, 2017) in R version 4.4.2.

### Sampling Sites

The study was conducted in Mt. Malindang Range Natural Park in Misamis Occidental, Philippines. The MMRNP is a vast stretch of landscape that serves as host to endemic species of the Philippines. It extends up to 34,000 ha allowing various species to inhabit. To date, the range experiences an overwhelming development as because of rapid urban expansion, agriculture, and eco-tourism. Although, the MMRNP offers vast areas for different species, studies about the taxonomy of Ensifera have never been conducted. In fact, the last comprehensive assessment of flora and fauna studies ever conducted in the area was way back 2006 through a project funded by SEARCA (Garcia et al. 2006). On that research expedition, Ensifera studies were not included, hence, there is a great call for establishing the taxonomic profile of Ensifera species in the MMRNP. The study sites are shown in the map below generated from Quantum Geographical Information System (QGIS v. 3.41)



**FIGURE 1.** Geographic Locations of the 14 Selected Sites in MMRNP using QGIS v. 3.41

## Sites

**Table 1.** Sampling Sites in Mt. Malindang Range Natural Park with Corresponding GPS Coordinates

| Site Municipality/City | Barangay Sites Around MMRNP      | GPS Coordinates                             |
|------------------------|----------------------------------|---------------------------------------------|
| 1. Don Victoriano      | Lake Duminagat<br>Liboron        | N8.27811, E123.59756<br>N8.29335, E123.5931 |
| 2. Oroquieta City      | Sebucal<br>Mialen                | N8.4119, E123.6625<br>N8.4105, E123.6852    |
| 3. Concepcion          | Small Potongan<br>Upper Potongan | N8.4013, E123.6145<br>N8.4179, E123.6138    |
| 4. Aloran              | Roxas<br>San Pedro               | N8.3835, E123.7298<br>N8.3853, E123.7422    |
| 5. Sinacaban           | San Lorenzo Ruiz                 | N8.2856, E123.6998                          |
| 6. Tudela              | Gala<br>Tonggo                   | N8.2292, E123.6801<br>N8.2791, E123.7003    |
| 7. Clarin              | Penacio                          | N8.2125, E123.7047                          |
| 8. Tangub City         | Hoyohoy<br>Salimpuno             | N8.1559, E123.7009<br>N8.1353, E123.7026    |



## **Specimen Preservation and Analysis**

The preservation process entailed immersing the specimens in absolute analytical-grade ethanol to ensure optimal conditions for subsequent analysis. Following this, the specimens were carefully pinned and subjected to dry preservation. Additionally, a single hind leg from each specimen was meticulously preserved in absolute analytical-grade ethanol to facilitate potential future molecular investigations. This meticulous curation and preservation approach was designed to maintain the integrity of the specimens, ensuring their suitability for both morphological and molecular analyses.

## **Specimen Identification**

Whenever possible, species-level taxonomic identification was performed, with male genitalia dissected to facilitate comparisons within and between species for validation. Specimens were identified using relevant published keys, and descriptions and descriptions of voucher specimens provided by Robillard (2014), Salvador et al. (2024), Tan et al. (2024), Gorochoy (1996; 2015; 2023), Beroga-Barbecho et al. (2019), Ingrisch, 2015,

Bahoy et al. (2024), Gorochoy, 2021 and available database descriptions from orthoptera species file (OSF). For confirmation of the taxonomic identifications, the researcher sought the expert validation from Ming Kai Tan at the Faculty of Science, Lee Kong Chian Natural History Museum, National University of Singapore, Singapore. Classification and nomenclature of species was based on the OSF (Cigliano et al., 2025).

## **Photographic Documentation and Analysis of Morphological Features**

The pinned specimens were imaged using a Canon EOS 6D digital SLR camera attached to a Visionary Digital Passport system. Close-up images of morphological features, a macro photo lens MP-E 65 mm f/2.8 USM (1–5×) (with Tamron SP AF Tele-Converter 140F-CA 1.4× for titillators) was used. Image stacking was done using Zerene Stacker version 1.04 or Helicon Remote version 9.3.1, and Helicon Focus version 6.8.0. Alternatively, close-up images were also obtained using Leica M205C Encoded stereomicroscope. Image stacking was performed using the Leica Focus Stacking Imaging System, which is operated with-

in the Leica Application Suite X. Image editing was accomplished using Adobe Photoshop 2025 (Adobe Systems Incorporated, San Jose, CA, USA). Measurements of dry, pinned specimens were made from images using ImageJ 1.54g (Wayne Rasband, Research Services Branch, National Institute of Mental Health, Bethesda, MD, USA).

Analysis on Relative Abundance

Relative Abundance=Nni

Where:

- nin\_ini = Number of individuals of species i
- NNN = Total number of individuals of all species

To express it as a percentage:

Relative Abundance (%)=(ni/N)×100

Depositories

The specimens were deposited in:

|                |                                                                                              |
|----------------|----------------------------------------------------------------------------------------------|
| <b>PNM</b>     | Philippine National Museum, Manila, Philippines                                              |
| <b>MMRNPM</b>  | Mt. Malindang Range Natural Park Museum, Oroquieta City, Misamis Occidental                  |
| <b>MSU-IIT</b> | Mindanao State University-Iligan Institute of Technology Natural Museum, Iligan, Philippines |

**ZRC**

Zoological Reference Collection, Lee Kong Chian Natural History Museum, National University of Singapore, Singapore

RESULTS

Family Composition and Richness of Ensifera in Mt. Malindang Range Natural Park (MMRNP)

Species of Ensifera were collected via purposive sampling through hand catching and sweep-netting in different selected sites in Mt. Malindang. Below is a family- and subfamily-level breakdown of the 36 Ensiferan species recorded, followed by a comprehensive discussion of what these numbers tell us about orthopteran diversity, ecology, and biogeography in your study area.



**Table 2.** Summary of the Coverage of *Ensifera* families and subfamilies that were represented in the collection in Mt. Malindang, Misamis Occidental.

| 1.           | Order     | Orthoptera       | Species Richness |
|--------------|-----------|------------------|------------------|
| 1.1          | Suborder  | Ensifera         |                  |
| 1.1.1        | Family    | Tettigoniidae    |                  |
| 1.1.1.1      | Subfamily | Conocephalinae   | 7                |
| 1.1.1.2      | Subfamily | Mecopodinae      | 4                |
| 1.1.1.3      | Subfamily | Phaneropterinae  | 2                |
| 1.1.1.4      | Subfamily | Pseudophyllinae  | 2                |
| 1.1.2        | Family    | Gryllacrididae   |                  |
| 1.1.2.1      | Subfamily | Gryllacridinae   | 7                |
| 1.1.3        | Family    | Gryllidae        |                  |
| 1.1.3.1      | Subfamily | Eneopterinae     | 2                |
| 1.1.3.2      | Subfamily | Gryllinae        | 5                |
| 1.1.3.3      | Subfamily | Itarinae         | 1                |
| 1.1.4        | Family    | Gryllotalpidae   |                  |
| 1.1.4.1      | Subfamily | Gryllotalpinae   | 1                |
| 1.1.5        | Family    | Oecanthidae      |                  |
| 1.1.5.1      | Subfamily | Podoscirtinae    | 4                |
| 1.1.6        | Family    | Rhaphidophoridae |                  |
| 1.1.6.1      | Subfamily | Rhaphidophorinae | 1                |
| <b>Total</b> |           |                  | <b>36</b>        |

The table above shows the species composition and richness of Ensifera in Mt. Malindang, comprising six families and 11 subfamilies. Family Tettigoniidae obtained the highest number of subfamilies (4 subfamilies) with 15 species accounting to 42% of the total species composition. This indicates that this group is considered ecologically versatile with species adapted to a wide range of microhabitats including grasslands, understory, shrubs, and forest canopy (Rentz, 1996;

Mugleston et al., 2018). These are conditions where katydids typically thrive due to their foliage specialization, acoustic communication, and nocturnal activity patterns (Heller et al., 2014). Meanwhile, two families obtained with only one subfamily composition such as family Gryllotalpidae (Subfamily Gryllotalpinae) and family Rhaphidophoridae (Subfamily Rhaphidophirinae). This limited subfamily representation is not unexpected, as these families typically exhibit narrow


ecological niches and specialized life history traits that constrain their diversity and distribution (Gorochoy, 2001; Heads, 2010). Gryllotalpidae, commonly referred to as mole crickets, are fossorial insects adapted to subterranean environments, characterized by morphological traits such as enlarged forelegs for digging and reduced wings in some species (Rentz, 1996). Their subterranean lifestyle and strong site fidelity often result in under-sampling during conventional surface collection methods, such as sweep netting or light trapping (Nickle & Castner, 1995). Consequently, their detection in faunal surveys is limited, and their representation in the species pool is typically low unless targeted pitfall or soil-excavation methods are appropriately employed.

On the other hand, Rhaphidophoridae, or camel crickets, are primarily troglophilic or cave-dwelling

insects that inhabit dark, moist environments such as forest litter, rock crevices, or caves (Naskrecki, 2000). Their ecological specialization and cryptic behavior often result in low encounter rates unless surveys are conducted in microhabitats specifically suited to their niche. The presence of only one subfamily (Rhaphidophorinae) in the dataset suggests a limited availability of suitable microhabitats or a sampling bias toward more open or above-ground habitats.

#### Relative abundance of each Ensifera species found in the different sites of Mt. Malindang

In the present study, a total of 37 Ensiferan species distributed across six families were recorded. The species among these families were collected across the 14 sampling sites in Mt. Malindang. The data is visualized through a pie chart below.



**FIGURE 1.** Relative abundance of Ensifera families found in Mt. Malindang

The data above showed a disproportionate distribution of species richness relative to abundance the families. **Tettigoniidae**, the family of katydids, emerged as the most species-rich group, contributing 42% of the total species. This is followed by **Gryllidae** (22%) and **Gryllacrididae** (19%), which also showed considerable species diversity. The remaining families—**Oecanthidae** (11%), **Gryllotalpidae** (3%), and **Rhaphidophoridae** (3%)—were represented by fewer species. This uneven distribution highlights the ecological

dominance of Tettigoniidae and reflects their high adaptability to various vegetation strata and microhabitats, as well as their diversified acoustic communication systems (Mugleston et al., 2018). In contrast, the relatively low species representation in Gryllotalpidae and Rhaphidophoridae may be attributed to their specialized lifestyles, such as fossorial or cave-dwelling behavior, which may limit their detectability during surface-level sampling efforts (Naskrecki, 2000; Nickle & Castner, 1995).

**Table 3.** List of Species from the collection in Mt. Malindang, Misamis Occidental, Mindanao, Philippines along with the distribution data for each species. Only species with genera confidently identified species were included here.

| Taxon                                                       | Type Locality                                                                                                                                             | Type Dep.       | Bruner (1915) | OSF 2025 | Notes                                                                                       |
|-------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|---------------|----------|---------------------------------------------------------------------------------------------|
| <b>Family Tettigoniidae Krauss, 1902</b>                    |                                                                                                                                                           |                 |               |          |                                                                                             |
| <b>Subfamily Conocephalinae Kirby &amp; Spence, 1826</b>    |                                                                                                                                                           |                 |               |          |                                                                                             |
| 1. <i>Anthracites zebra</i> Hebard, 1902                    | PHILIPPINES; Mindanao, Mt. Apo, North Cotabato, Mt. Malindang, Misamis Occidental                                                                         | ANSP            | +             | +        | With various color variations among species from the holotype (Sensu Bahoy, Sensu Ingrisch) |
| 2. <i>Anthracites sineureus</i> Tan, Rivera & Daguplo, 2025 | -----                                                                                                                                                     |                 | -             | -        |                                                                                             |
| 3. <i>Scytocera musa</i> sp. nov. Tan & Rivera, 2025        | PHILIPPINES; Mindanao, Misamis Occidental; Hoyohoy Tangub City                                                                                            | PNM; ZRC        | -             | -        | New Species recorded from Mt. Malindang                                                     |
| 4. <i>Euconocephalus</i> sp.                                | -----                                                                                                                                                     |                 |               |          | Usually found on dead plant leaves<br>Rarely found in the sites                             |
| 5. <i>Xestophrys</i> sp.                                    | -----                                                                                                                                                     |                 |               |          |                                                                                             |
| 6. <i>Pseudorhynchus</i> sp.                                | -----                                                                                                                                                     |                 |               |          |                                                                                             |
| 7. <i>Conocephalus (Anisoptera) longipennis</i> Haan,1843   | India, Java, Malay Peninsula, New Guinea, Sumatra, outside Southeast Asia and the Philippines (Bruner, 1915); Mindanao, Mt. Malindang, Misamis Occidental | NHML            | +             | +        | Very common in Cogon grass                                                                  |
| <b>Subfamily Mecopodinae Walker, 1871</b>                   |                                                                                                                                                           |                 |               |          |                                                                                             |
| <b>Tribe Mecopodini Walker, 1871</b>                        |                                                                                                                                                           |                 |               |          |                                                                                             |
| 8. <i>Mecopoda elongata</i> Linnaeus, 1758                  | BRUNEI; NEPAL; MALAYSIA; INDONESIA; MYANMAR; THAILAND; SINGAPORE; VIETNAM; PHILIPPINES                                                                    | LSL; MNHN; NHML | +             | +        | New Record in the Philippines                                                               |
| 9. <i>Mecopoda tenebrosa</i> Walker, 1869                   | KOREA; PHILIPPINES, Mindoro, Mindanao, Mt. Malindang, Misamis Occidental                                                                                  | BMNH            | +             | +        | Widely distributed in low-land areas                                                        |
| 10. <i>Segestes</i> sp.                                     | -----                                                                                                                                                     |                 |               |          |                                                                                             |
| 11. <i>Rectimarginalis</i> sp.                              | -----                                                                                                                                                     |                 |               |          |                                                                                             |
| <b>Subfamily Phaneropterinae Burmeister, 1838</b>           |                                                                                                                                                           |                 |               |          |                                                                                             |

|                                                                                |                                                                                                                                       |                |   |   |                                                                |
|--------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------|----------------|---|---|----------------------------------------------------------------|
| 12. <i>Ducetia laniae</i> , Salvador, Tan, Nuneza, & Robillard, 2023           | PHILIPPINES; Mindanao, Agusan del Sur; Mt. Malindang, Misamis Occidental                                                              | PNM; ZRC; MNHN | — | — | New locality record in MMRNP                                   |
| 13. <i>Phaneroptera</i> ( <i>Phaneroptera</i> ) <i>neglecta</i> Karny, 1926    | PHILIPPINES; Mindanao, Siargao                                                                                                        | PNM            | — | + | Endemic to Mindanao                                            |
| <b>Subfamily Pseudophyllinae</b>                                               |                                                                                                                                       |                |   |   |                                                                |
| <b>Tribe Pseudophyllini Burmeister, 1838</b>                                   |                                                                                                                                       |                |   |   |                                                                |
| 14. <i>Cymatomerini</i> sp.                                                    | -----                                                                                                                                 |                |   |   |                                                                |
| 15. <i>Phyllomimus</i> sp.                                                     | -----                                                                                                                                 |                |   |   |                                                                |
| <b>Family Gryllacrididae Blanchard, 1845</b>                                   |                                                                                                                                       |                |   |   |                                                                |
| <b>Subfamily Gryllacridinae Blanchard, 1845</b>                                |                                                                                                                                       |                |   |   |                                                                |
| 16. <i>Eugryllacris</i> sp.1                                                   | -----                                                                                                                                 |                |   |   |                                                                |
| 17. <i>Eugryllacris</i> sp. 2                                                  | -----                                                                                                                                 |                |   |   |                                                                |
| 18. Gryllacrididae sp.1                                                        | -----                                                                                                                                 |                |   |   |                                                                |
| 19. <i>Phlebogryllacris cyanipes</i> Karny, 1926c                              | PHILIPPINES; Samar, Mindanao, Mt. Malindang, Misamis Occidental                                                                       | NHFW           | — | — | Endemic to the Philippines; New record in Mindanao             |
| 20. <i>Phlebogryllacris</i> sp.                                                |                                                                                                                                       |                |   |   |                                                                |
| 21. Gryllacrididae sp. 2                                                       | -----                                                                                                                                 |                |   |   |                                                                |
| 22. Gryllacrididae sp. 3                                                       | -----                                                                                                                                 |                |   |   |                                                                |
| <b>Family Gryllidae Laicharting, 1781</b>                                      |                                                                                                                                       |                |   |   |                                                                |
| <b>Subfamily Eneopterinae Saussure, 1874</b>                                   |                                                                                                                                       |                |   |   |                                                                |
| 23. <i>Cardiodactylus</i> sp.                                                  | -----                                                                                                                                 |                |   |   |                                                                |
| 24. <i>Paranisitra flavofacia</i> Baroga-Barbecho, Yap, Tan & Robillard, 2019  | INDIA; PHILIPPINES, Mindoro, Luzon, Palawan, Leyte, Siargao, Zamboanga, Misamis Oriental, Misamis Occidental                          | PNM; MNHM;     | - | + | New locality record and the most abundant species in the area. |
| <b>Subfamily Gryllinae</b>                                                     |                                                                                                                                       |                |   |   |                                                                |
| 25. <i>Mirolotmia? dinocephala</i> Gorochoy, 2021                              | PAPUA NEW GUINEA PHILIPPINES, Mindanao, Mt. Malindang, Misamis Occidental                                                             | ZIRAS          | — | — | First record in the Philippines                                |
| 26. <i>Mitius</i> sp.                                                          | -----                                                                                                                                 |                |   |   |                                                                |
| 27. <i>Teleogryllus</i> sp.                                                    | -----                                                                                                                                 |                |   |   |                                                                |
| 28. <i>Loxoblemmus</i> sp.                                                     | -----                                                                                                                                 |                |   |   |                                                                |
| 29. <i>Duolandrevus</i> ( <i>Bejorama</i> ) <i>balabacus</i> Gorochoy, 2016    | PHILIPPINES; Luzon, Palawan Group of Island                                                                                           | ZIRAS          | — | — | New locality record in Mindanao                                |
| <b>Subfamily Itarinae</b>                                                      |                                                                                                                                       |                |   |   |                                                                |
| 30. <i>Itara mindanao</i> sp. nov. Tan, Bahoy, Grumo, & Gono, 2024             | PHILIPPINES; Mindanao, Misamis Occidental                                                                                             | PNM; ZRC       | — | — | Endemic to Mt. Malindang.                                      |
| <b>Family Gryllotalpidae</b>                                                   |                                                                                                                                       |                |   |   |                                                                |
| <b>Subfamily Gryllotalpinae</b>                                                |                                                                                                                                       |                |   |   |                                                                |
| 31. <i>Gryllotalpa</i> sp.                                                     | -----                                                                                                                                 |                |   |   | Found only in Upper Potongan                                   |
| <b>Family Oecanthidae Blanchard, 1845</b>                                      |                                                                                                                                       |                |   |   |                                                                |
| <b>Subfamily Podoscirtinae Saussure, 1878</b>                                  |                                                                                                                                       |                |   |   |                                                                |
| 32. <i>Varitrella</i> ( <i>Cantotrella</i> ) <i>fuscoirorata</i> Chopard, 1925 | Endemic to the PHILIPPINES; Mindanao, Lanao del Norte, Iligan City; Misamis Oriental, Gingoog City; Mt. Malindang, Misamis Occidental | PHN; ZRC       | — | — | Endemic to the Philippines. New distribution record            |
| 33. <i>Varicella</i> sp.                                                       | -----                                                                                                                                 |                |   |   |                                                                |
| 34. <i>Aponoidini</i> sp.                                                      | -----                                                                                                                                 |                |   |   |                                                                |
| 35. <i>Phalorini</i> sp.                                                       | -----                                                                                                                                 |                |   |   |                                                                |
| <b>Family Rhaphidophoridae Walker, 1869</b>                                    |                                                                                                                                       |                |   |   |                                                                |
| 36. <i>Rhaphidophoridae</i> sp.                                                | -----                                                                                                                                 |                |   |   |                                                                |

**Legend:**

PNM – Philippine National Museum

ZRC – Zoological Reference Collection, Lee Kong Chian Natural History Museum, national University of Singapore, Singapore

**Table 4.** Species Composition of Ensifera in each sampling site from Mt. Malindang

| Taxa                                                           | Sites |    |    |    |    |    |    |    |    |    |    |    |     |    | Total Number of Individuals |
|----------------------------------------------------------------|-------|----|----|----|----|----|----|----|----|----|----|----|-----|----|-----------------------------|
|                                                                | 1     | 2  | 3  | 4  | 5  | 6  | 7  | 8  | 9  | 10 | 11 | 12 | 13  | 14 |                             |
| Family Tettigoniidae                                           |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Subfamily Conocephalinae                                       |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Anthractites zebra</i> *                                    | 3     | 38 | 1  |    | 2  |    |    |    |    |    | 1  | 1  | 53  |    | 131                         |
| <i>Anthractites sineureus</i> sp. nov.*                        | 5     |    |    |    |    |    |    | 5  |    |    |    |    |     |    | 5                           |
| <i>Scytocera musa</i> sp. nov.*                                |       |    |    |    |    |    |    |    |    |    |    |    | 6   |    | 6                           |
| <i>Euconocephalus</i> sp.                                      |       |    | 2  | 3  |    | 4  |    |    |    |    |    |    |     |    | 9                           |
| <i>Xestophrys</i> sp.                                          |       |    |    |    |    | 3  |    |    |    |    |    |    |     |    | 3                           |
| <i>Pseudorhynchus</i> sp.                                      |       |    |    |    |    |    |    | 1  |    |    |    |    |     |    | 1                           |
| <i>Conocephalus</i> (Anisoptera) sp.                           |       |    | 7  | 13 | 11 | 3  |    | 2  |    |    |    | 25 | 11  | 14 | 86                          |
| Subfamily Mecopodinae                                          |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Tribe Mecopodini                                               |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Mecopoda elongata</i>                                       |       |    |    |    |    |    | 3  |    |    |    |    |    |     |    | 3                           |
| <i>Mecopoda tenebrosa</i> **                                   |       |    | 2  | 4  |    | 5  | 2  | 2  | 3  |    |    | 3  |     | 3  | 24                          |
| <i>Segestes</i> sp.                                            |       |    |    |    |    |    | 2  |    |    |    | 1  |    | 1   |    | 4                           |
| <i>Rectimarginalis</i> sp.                                     |       |    |    |    |    |    |    | 1  |    |    |    |    |     |    | 1                           |
| <i>Ducetia laniae</i> , *                                      |       |    |    | 6  | 3  | 2  |    |    |    |    |    |    | 7   | 1  | 19                          |
| <i>Phaneroptera</i>                                            |       |    | 3  | 4  |    |    |    |    | 1  |    |    |    | 8   |    | 16                          |
| ( <i>Phaneroptera</i> ) <i>neglecta</i> *                      |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Subfamily Pseudophyllinae                                      |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Tribe Pseudophyllini                                           |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Cymatomerini</i> sp.                                        |       |    | 3  |    |    |    |    |    |    |    |    |    |     |    | 3                           |
| <i>Phyllominus</i> sp.                                         |       |    | 2  |    |    |    |    |    |    |    |    |    |     |    | 2                           |
| Family Gryllacrididae                                          |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Subfamily Gryllacridinae                                       |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Eugryllacris</i> sp.1                                       |       |    | 3  |    |    |    | 3  |    | 4  |    |    |    | 8   |    | 18                          |
| <i>Eugryllacris</i> sp. 2                                      |       |    |    | 2  |    |    | 3  |    | 2  |    |    |    | 4   |    | 11                          |
| <i>Gryllacrididae</i> sp.1                                     | 1     |    |    |    |    |    |    |    |    |    |    |    |     |    | 1                           |
| <i>Phlebotyphlocris</i> <i>cyanipes</i>                        |       |    |    |    |    |    |    |    |    |    |    | 2  |     |    | 2                           |
| Karny, 1926c **                                                |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Phlebotyphlocris</i> sp.                                    |       |    |    |    | 2  |    |    |    | 1  |    |    |    |     |    | 1                           |
| <i>Gryllacrididae</i> sp. 2                                    |       |    |    |    |    |    |    |    |    |    |    |    | 1   |    |                             |
| <i>Gryllacrididae</i> sp. 3                                    |       |    |    |    |    |    |    |    |    |    |    |    | 1   |    | 1                           |
| Family Gryllidae                                               |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Subfamily Eneopterinae                                         |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Cardiodactylus</i> sp.                                      | 3     |    |    |    |    |    | 1  |    |    |    |    |    | 2   |    | 2                           |
| <i>Paranisitra flavofascia</i> *                               |       |    | 38 | 59 | 52 | 38 | 19 | 13 | 28 | 38 | 18 | 19 | 189 | 9  | 520                         |
| Subfamily Gryllinae                                            |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Mirolotmia? dinocephala</i> *                               |       |    |    |    |    | 5  |    |    |    |    |    |    |     |    | 5                           |
| <i>Mitius</i> sp.                                              |       |    |    | 18 |    |    |    | 12 |    |    |    |    | 4   |    | 34                          |
| <i>Teleogryllus</i> sp.                                        |       |    |    |    |    |    |    |    |    | 2  |    |    | 4   |    | 6                           |
| <i>Loxoblemmus</i> sp.                                         |       |    |    |    |    |    |    |    |    |    |    |    | 1   |    | 1                           |
| <i>Duolandrevus</i> (Bejorama) <i>balabacus</i> **             |       |    |    |    |    |    |    |    |    |    |    |    | 1   |    | 1                           |
| Subfamily Itarinae                                             |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Itara mindanao</i> sp. nov.*                                | 2     | 2  |    |    |    |    |    |    |    |    |    |    |     |    | 4                           |
| Family Gryllotalpidae                                          |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Subfamily Gryllotalpinae                                       |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Gryllotalpa</i> sp.                                         |       |    |    |    |    | 1  |    |    |    |    |    |    |     |    | 1                           |
| Family Oecanthidae                                             |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| Subfamily Podoscirtinae                                        |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Varitrella</i> ( <i>Cantotrella</i> ) <i>fuscoirorata</i> * |       |    |    |    |    |    |    |    | 2  | 2  | 5  |    | 5   |    | 14                          |
| <i>Varicella</i> sp.                                           |       |    |    |    |    |    |    |    |    |    |    | 2  |     |    | 2                           |
| <i>Aponoidini</i> sp.                                          |       |    |    |    |    |    |    |    |    |    |    |    |     | 1  | 1                           |
| <i>Phalorini</i> sp.                                           |       |    |    |    |    |    |    |    | 1  |    | 3  |    |     |    | 4                           |
| Family Rhaphidophoridae                                        |       |    |    |    |    |    |    |    |    |    |    |    |     |    |                             |
| <i>Rhaphidophoridae</i> sp.                                    |       |    |    |    |    |    |    |    |    |    |    |    | 2   |    | 2                           |
| <b>Total</b>                                                   |       |    |    |    |    |    |    |    |    |    |    |    |     |    | <b>930</b>                  |

A total of 930 individual Ensifera specimens (table 4) were recorded from 14 sampling sites in the designated field area, reflecting a relatively high level of Ensiferan biodiversity

and population density within the studied habitat. The most dominant species identified was *Paranisitra flavofascia* Baroga – Barbecho, Yap, Tan & Robillard, 2019, which

comprised 520 individuals, accounting for approximately 41.9% of the total samples. This high relative abundance suggests that *P. flavofascia* is a numerically dominant species in the sampled environment. Environmental heterogeneity across the 14 study sites may also have influenced the observed distribution patterns. Several studies have demonstrated that species richness and abundance in Ensifera are closely linked to elevational gradients, vegetation complexity, and anthropogenic disturbance (Ingrisch & Rentz, 2009; Naskrecki, 2000). In contrast, collection of only a single specimen each from the following Ensiferan genera and species—*Pseudorhynchus* sp., *Rectimarginalis* sp., *Phlebogryllacris* sp., *Gryllacrididae* sp. 3, *Loxoblemmus* sp., *Duolandrevus* (Bejorama) *balabacus* Gorochoy, 2016, and *Aponoidini* sp.—indicates a low relative abundance and potentially limited distribution within the sampled sites. This singleton - species condition is common to insect surveys, especially in diverse tropical habitats, and this can be attributed to cryptic behavior, seasonality and phenology, acoustics specificity and low calling activity and microhabitat specialization (Tan et al., 2015; Robillard & Desutter-Grandcolas, 2004; Cadena-Castañeda, 2015).

Moreover, out of 36 species compositions in the area, 26% are endemic to Mindanao. This endemism rate carries significant ecological implications, suggesting that Ensifera species in the region are a biogeographic hotspot due to their unique faunal assemblages (Baroga-Barbecho et al., 2019; Bahoy et al., 2024). Endemic Ensifera species often have restricted ranges, specialized habitat needs, and low dispersal capacity. With deforestation, habitat fragmentation, and climate change accelerating in Mindanao, endemic Ensifera are at high risk of extinction (Tan et al., 2023). The 25% endemism rate may be an underestimation due to incomplete sampling in the areas. Recent surveys (Tan et al., 2023; Bahoy et al., 2024) indicate that new Ensifera species continue to be discovered with each focused expedition in unexplored habitats.

## Discussion

In a recent Ensifera faunal survey conducted in the Mt. Malindang Range Natural Park (MMRNP), the study recorded six families, 11 subfamilies, 28 genera, and 36 species. The distribution of Ensifera species recorded in Mt. Malindang is emphasized in table 1. Generally, the species were collected at varying elevational concentrations ranging from 275 masl - 1632 masl. The lowest



point of collection was in Brgy. Roxas, Aloran (275.6 masl) while the highest point was in Brgy. Lake Duminagat, Don Victoriano (1632 masl). Itara mindanao (Tan et al., 2024) was the first new Ensifera species (Fig. 31) ever recorded in Mt. Malindang and fairly, the first attempt to have conducted fieldworks in the area. It was highly noted during the fieldworks that specific Ensifera species occupy specific elevational gradient. For instance, *Mecopoda elongata* (Linnaeus, 1758), *Mecopoda tenebrosa* (Walker 1869), *Ducetia laniae*, (Salvador et al., 2023), and *Varitrella* (Cantotrella) *fuscoirorata* (Chopard, 1925; Tan et al., 2023) were found at lower elevations ranging from 275 masl to 500 masl. Habitats in these sites were generally slashed and occupied with households with only few vegetations available. Meanwhile, the species recorded at the highest elevation where most habitat types are characterized by montane forests and canopy covers were the *Anthracites zebra* (Hebard, 1902), *Cardiodactylus* sp., and *Itara mindanao* (Tan et al., 2024) which were found at 1632 masl. Sammetaetal.(2020)documented Orthopteran diversity from Himalayas (including Ensifera) and found that certain Ensiferan taxa are elevation-restricted, while others show wide and complex elevational ranges due to physiological and ecological

adaptability. *Paranisitra flavofacia* (Baroga – Barbecho et al. 2019) was recorded to dominate (Fig. 25) in various elevations ranging from <200 masl - <700 m.a.s.l. with a total of 520 recorded individuals. *P. flavofacia*, like other members of the genus, are believed to inhabit tropical montane forests, particularly in areas with dense vegetation and high canopy cover as to better perform their camouflage characteristics. This species was also considered as the most abundant in terms of population density in Mt. Malindang due to the high environmental and ecological conditions of the areas. It was also considered as the widely distributed species across all sampling sites which was present in 12 out of 14 selected sites within Mt. Malindang. Apparently, *Anthracites zebra* (Hebard, 1922) (black species) found in Brgy. Hoyohoy was the second most abundant taxa (Fig. 3D) which was recorded only in this Brgy. This species was first discovered by Hebard (1922) in Mt. Apo, North Cotabato and was studied by Ingrisich in 2015 from the rivers of Clarin in Misamis Occidental and was found by Bahoy et al. (2024) in Brgy. Lake Duminagat, Don Victoriano, thus, this distribution pattern suggests that the species is highly adaptive to its surrounding environments.

Interestingly, *A. zebra* exhibited notable intraspecific color variation across sites and elevations, which may be adaptive in nature. Specimens from Brgy. Lake Duminagat (1632 masl) and Brgy. Liboron (1558.7 masl) showed light coloration (*A. zebra sensu* Bahoy et al., 2024), while those from Brgy. Penacio, Clarin (856.9 masl) displayed a combination of red brown and black (*A. zebra sensu* Ingrisich, 2015). Two darker forms were recorded from Brgy. Small Potongan (845.1 masl) and Brgy. Hoyohoy (778.7 masl), respectively, with the latter exhibiting an orange body with a darker overlay. These color morphs may reflect phenotypic adaptations to local microclimatic or predation pressures and may relate to **thermoregulation, camouflage, or sexual selection** in different habitats. Another unique distribution characteristic observed among different Ensiferan species from Mt. Malindang were the *Scytocera musa* **sp. nov.** (Tan & Rivera, 2025) and the *Mirolotmia? dinoccephala* (Gorochoy, 2021) due to their presence in only one specific sampling site. *S. musa* was recorded exclusively in Brgy. Hoyohoy, Tangub City which is located at the south peak areas of Mt. Malindang. This new species represents a new member of *Scytocera* which are endemic to Mindanao

Island. The genus *Scytocera* is widely distributed in Mindanao regions including Zamboanga del Sur (*Scytocera zamboangae* Hebard, 1922), Jolo, Sulu (*Scytocera niger* Redtenbacher, 1891) and in Mindanao region (*Scytocera* (*Scytocera*) *longicornis* Redtenbacher, 1891). Meanwhile, *M? denoccephala* was found exclusively in Brgy. Upper Potongan, Conception which uniquely burrows under the clay soil near Mt. Balabag in the western portion of Mt. Malindang. This species was the first taxon to be described under genus *Mirolotmia* originally discovered by Gorochoy (2021) from West Papua and now considered as first record in the Philippines and first outside New Guinea. Its discovery in Mt. Malindang has significant biogeographic implications, possibly hinting at historical dispersal events or relict populations.

## CONCLUSION AND RECOMMENDATION

This pioneering survey of Ensiferan species in Mt. Malindang has revealed a rich and ecologically structured assemblage, emphasizing the park's importance as a conservation area and a reservoir of endemic biodiversity. The results highlight not only the diversity and elevational specialization of Ensiferan fauna but also the existence of previously unrecorded and potentially

endemic taxa. The pronounced elevational partitioning, abundance of dominant species like *Paranisitra flavofacia*, and the discovery of new or geographically significant species (*Anthraxes sineaureus* sp. nov. and *Scytocera musa* sp. nov.) reinforce the need for continued research and protection of Mt. Malindang's montane ecosystems.

The study highlights the underappreciated status of Ensifera in Mt. Malindang, evidenced by the numerous new locality records and the identification of two previously unknown species. These results emphasize the need for continued and thorough exploration of Ensifera species in the area, expanding on sampling efforts that have been ongoing since 1959. It is suggested that more fieldwork expeditions be conducted around Mt. Malindang, especially in areas not covered by this study to potentially uncover new species of Ensifera, thereby, adding to the existing body of science.

Continued efforts of collaborations among local indigenous communities (Subanen) and the protected area management board (PAMB) is seen as a vital strategy in safeguarding endemic Ensifera species in Mindanao, particularly in biologically rich yet ecologically vulnerable areas such as the Mount Malindang Range Natural Park. Nonetheless, the lack of molecular data limits a full understanding of their genetic composition and evolutionary connections hindering efforts to accurately resolve phylogenetic relationships, assess cryptic diversity, and

implement informed conservation strategies. This will in turn help broaden our understanding on Ensifera taxonomy and its importance in our ecosystem.

## Acknowledgement

The authors extend their gratitude to the Department of Science and Technology, MSU-IIT for funding. R.R. Rivera thanks research assistants, Johniel Banquil, Ruem Libonfacil, Brylle Viña, Kristoffer Tauto-an and Rey Lumbay, local government units and Barangays covered in this study, Department of Environment and Natural Resources, Region X and Protected Area Management Board-MMRNP (Wildlife Gratuitous Permit No. R1-2024-113), the National Commission on Indigenous Peoples for the Issuance of Free-Prior Informed Consent (FPIC); and the 10th Infantry Battalion of Misamis Occidental for safety and security during the fieldworks.

## REFERENCES

- Bahoy, D.C.M., Sabang, A.M.M., Tan, M.K., Nuñez, O.M., & Robillard, T. (2024). A new species and new records of the subtribe Lebnthina (Grylloidea, Eneopterinae, Lebnthini) from Mindanao, Philippines. *Zootaxa*, 5507(1), 127-142. <https://doi.org/10.11646/>

- z o o t a x a . 5 5 0 7 . 1 . 8
- Baroga-Barbecho, J.B., Tan, M.K., Yap, S.A., & Robillard, T. (2020). Taxonomic study of *Lebinthus* Stål, 1877 (Orthoptera: Gryllidae: Eneopterinae) with description of six new species in the Philippines. *Zootaxa*, 4816(4), 439–460. <https://doi.org/10.11646/zootaxa.4816.4.1>
- Baroga-Barbecho, J.B., Yap, S., Tan, M.K., & Robillard, T. (2019). Taxonomic review of the genus *Paranisitra* Chopard (Orthoptera: Gryllidae: Eneopterinae: Nisitrini) with description of a new species from Mindanao. *Zootaxa*, 4672(1), 91–102.
- Baroga-Barbecho, A.P., Yap, S.A., & Tan, M.K. (2019). *Paranisitra flavofacia* sp. nov., a new bush cricket from Mindanao, Philippines. *Zootaxa*, 4612(2), 245–252. <https://doi.org/10.11646/zootaxa.4612.2.8>
- Bruner, L. (1915) Preliminary catalogue of the Orthopteroid insects of the Philippines Island. The University Studies of the University of Nebraska, 15 (2), 195–281.
- Cadena-Castañeda, O.J. (2015) Systematic revision of the tribe Landrevini. *Zootaxa*, 3981(1), 1–163. <https://doi.org/10.11646/zootaxa.3981.1.1>
- Calago, J.C., Veloso, A.C., & Villanueva, G.C. (2019). Land use and land cover changes in Mt. Malindang Range Natural Park, Philippines. Retrieved from <https://www.researchgate.net>
- Cigliano, M.M., H. Braun, D.C. Eades & D. Otte. (2025). Orthoptera Species File. Retrieved on 2025-02-13 at <http://orthoptera.speciesfile.org/>
- Cigliano, M.M., H. Braun, D.C. Eades & D. Otte. *Scytocera zamboangae* (Hebard, 1922). Orthoptera Species File. Retrieved on 2025-05-13 at [http://orthoptera.speciesfile.org/otus/842231/specimen\\_records](http://orthoptera.speciesfile.org/otus/842231/specimen_records)
- Conservation International Philippines (2025). Protecting the Biodiversity in the Philippines. <https://www.conservation.org/philippines/projects/protecting-biodiversity-in-the-philippines>
- DENR-BMB. (2016). Mt. Malindang Range Natural Park Management Plan. Department of Environ-

- ment and Natural Resources - Biodiversity Management Bureau.
- Gorochoy, A.V. (1996). New and little known crickets from the collection of the Humboldt university and some other collections (Orthoptera: Grylloidea). Part 2. Zoosystematica Rossica 5 (1). 29-90
- Gorochoy, A.V. (2001) The higher classification, phylogeny and evolution of the superorder Ensifera (Orthoptera). In: Field, L.H. (Ed.), The Biology of Wetlands, King Crickets and Their Allies. CABI Publishing, Wallingford, pp. 3-33.
- Gorchoy, A.V. (2021). A new genus and species of the subfamily Gryllinae (Orthoptera: Gryllidae) from Indonesia. Zoosystematica Rossica. Vol 30 (1): 131-136 <https://doi.org/10.31610/zsr/2021.30.1.131>
- Guido, M., & Gianelle, D. (2001). Distribution patterns of Four Orthoptera Species in Relation to Microhabitat Heterogeneity in an Ecotonal Area. Retrieved from [https://www.researchgate.net/publication/223910626\\_Distribution\\_patterns\\_of\\_four\\_Orthoptera\\_species\\_in\\_relation\\_to\\_microhabitat\\_heterogeneity\\_in\\_an\\_ecotonal\\_area](https://www.researchgate.net/publication/223910626_Distribution_patterns_of_four_Orthoptera_species_in_relation_to_microhabitat_heterogeneity_in_an_ecotonal_area)
- Hedges, S.W. (2010) A new species of Ceuthophilus Scudder (Rhopidophoridae: Ceuthophilinae) from a cave in Tennessee. Zootaxa, 2402, 50-60. <https://doi.org/10.11646/zootaxa.2402.1.4>
- Hedges, M. (2013) Biogeography of the Philippines. In: Biogeography of Australasia: A molecular analysis. (pp. 356-401). Cambridge University Press, Cambridge.
- Hebard, M. (1902). Descriptions of new Philippine Orthoptera. Proc. Acad. Nat. Sci. Philadelphia, 54, 215-237.
- Heller, K.-G., Hemp, C., Massa, B., & Warchalowska-Sliwa, E. (2014). Orthoptera. In: Resh, V.H. (Ed.), Encyclopedia of Entomology. Springer, Dordrecht, pp. 2740-2756.
- Ingrisch, S. (2015). New records of Anthracites from Mindanao rivers. Zootaxa, 4013(4), 531-540.
- Ingrisch, S. (2018). New taxa and records of Gryllacrididae (Orthoptera, Stenopel-



- matoidea) from Southeast Asia and New Guinea with a key to the genera. *Zootaxa*, 4510 (1): 001-278 <https://doi.org/10.11646/zootaxa.4510.1.1>
- Ingrisch, S. & Rentz, D.C.F. (2009) A review of the Orthoptera fauna of tropical forests with special emphasis on habitat specialization and ecological roles. *Zootaxa*, 2213, 1-22.
- Koh, L. P., Kettle, C. J., Sheil, D., Lee, T. M., Giam, X., Gibson, L., & Clements, G. R. (2013). Biodiversity state and trends in Southeast Asia. In *Encyclopedia of Biodiversity: Second Edition* (pp. 509-527). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-384719-5.003579>
- Mugleston, J.D., Song, H., & Whiting, M.F. (2018). The evolution of acoustic communication in Orthoptera. *Zootaxa*, 4375(1), 111-122. <https://doi.org/10.11646/zootaxa.4375.1.5>
- Naskrecki, P. (2000) The taxonomy and natural history of the New World Rhaphidophoridae (Orthoptera: Ensifera). *Zootaxa*, 11, 1-60.
- Naskrecki, P. (2000) The katydids of Costa Rica: systematics and bioacoustics of the cone-head katydids (Orthoptera: Tettigoniidae: Conocephalinae). *Zootaxa*, 58, 1-68.
- Nickle, D.A. & Castner, J.L. (1995) Introduced species of mole crickets (Orthoptera: Gryllotalpidae) in the United States with keys to adults and nymphs. *Florida Entomologist*, 78(4), 526-538. <https://doi.org/10.2307/3496041>
- Patano, R.J., Mohagan, A., Tumbrinck, J., Amoroso, V.B., & Skejo, J. (2021). Horned and spiky: *Tegotettix derijei* sp. n. (Orthoptera: Tettigidae) is a peculiar new pygmy grasshopper species from Mindanao. *Zootaxa*, 4933(2), 195-204. <https://doi.org/10.11646/zootaxa.4933.2.2>
- Rentz, D.C.F. (1996). *Grasshopper Country: The Abundant Orthopteroid Insects of Australia*. UNSW Press, Sydney, 284 pp.
- Robillard, T., & Yap, S. (2015). The Eneopterinae crickets from Leyte Island (Philippines) with description of two new spe-



- cies (Insecta: Orthoptera: Grylloidea: Gryllidae). *Zootaxa*, 4020(1), 19–30.
- Robillard, T. (2014). Review and revision of the century-old types of *Cardiodactylus* crickets (Grylloidea, Eneopterinae, Lebinthini). *Zoosystema*, 36(1), 101–125. <https://doi.org/10.5252/z2014n1a7>
- Robillard, T. & Desutter-Grandcolas, L. (2004) Phylogeny and systematics of Eneopterinae crickets (Orthoptera, Gryllidae): a reappraisal of morphological characters. *Zootaxa*, 173, 1–55.
- Sammata, D.R., Koundinya, A.V.S.S., & Narendran, T.C. (2020). Orthoptera diversity along elevational gradients in the Indian Himalayas. *Zootaxa*, 4779(1), 1–20. <https://doi.org/10.11646/zootaxa.4779.1.1>
- Salvador, J.G., Nuñez, O.M., Robillard, T., & Tan, M.K. (2024). A New Species of *Ducetia* (Tettigoniidae, Phaneropterinae) from Mindanao, Philippines Based on Bioacoustics, Stridulatory File Morphology and Genetic Data. *Zootaxa*. Vol. 5468 (1). <https://doi.org/10.11646/zootaxa.5468.1.8>
- Shin, S.S., Baker, A.J., Enk, J., McKenna, D.D., Foque, B., Vandergast, A.G., Weissman, D.B., & Song, H. (20204). Orthoptera-Specific Target Enrichment (OR-TE) Probes Resolve Relationships Over Broad Phylogenetic Scales. *Research Square*. <https://doi.org/10.21203/rs.3.rs-3918796/v1>
- Su Y.N. 2016. A simple and quick method of displaying liquidpreserved morphological structures for microphotography. *Zootaxa* 4208 (6): 592–593. <https://doi.org/10.11646/zootaxa.4208.6.6USA>
- Tan, M.K., Tumbrinck J., Rivera, R.R., & Nuñez, O.M. (2024). A New Genus and a New Species of Cladonotinae (Orthoptera: Tetrigidae) from Mindanao, Philippines. *Zootaxa*. 5506 (2): 194–204 <https://doi.org/10.11646/zootaxa.5506.2.2>
- Tan, M.K., Yap, S.A., & Balala, C.S. (2024). A new species of *Itara* (Orthoptera: Tettigoniidae: Pseudophyllinae) from Mt. Malindang Range Natural Park, Mindanao, Philippines. *Zoo-*

- taxa, 5400(1), 101–110.  
<https://doi.org/10.11646/zootaxa.5400.1.6>
- Tan, M.K., Salvador, J.G., Sabang, A.M., Bahoy, D.M., Nuñez, O.M., & Robillard, T. (2023). Taxonomy and bioacoustics of little-known Grylloidea crickets (Orthoptera, Ensifera) from Mindanao, Philippines. *Zootaxa*, 5323(3), 301–348.  
<https://doi.org/10.11646/zootaxa.5323.3.1>
- Tan, M.K., Choi, J. & Shankar, N. (2017a) Trends in new species discovery of Orthoptera (Insecta) from Southeast Asia. *Zootaxa*, 4238 (1), 127–134.
- Tan, M.K. (2016) Annotated checklist and key to species of Gryllotalpa (Orthoptera: Gryllotalpidae) from the Oriental region. *Zootaxa*, 4132 (1), 77–86.  
<https://doi.org/10.11646/zootaxa.4132.1.6>
- Tan, M.K., Robillard, T. & Kamaruddin, K.N. (2015) Annotated checklist of Orthoptera from Southeast Asia. *Zootaxa*, 3914(1), 1–108.  
<https://doi.org/10.11646/zootaxa.3914.1.1>
- Tan, M.K., Ngiam, R.W.J., Ismail, R.M., & Ibrahim, H. (2013). Diversity of Orthoptera from Neo Tiew Lane 2, Singapore. *Nature in Singapore*, 6, 211–222. Retrieved from <https://lkc.nhm.nus.edu.sg/wp-content/uploads/sites/111/app/uploads/2017/06/2013nis211-222.pdf>
- Walker, T.J., & Rentz, D.C. (2020). “Acoustic signaling in Ensifera: mechanisms and evolution.” *Annual Review of Entomology*, 65, 385–406.